

to the apex; the apical margin well defined and crenulate, extending laterally halfway up the declivity.

Length, 2.1 mm.; breadth, .9 mm.

Bengal: Raja Bhat Khawa (*C. F. C. Beeson*), ex Sal twig.

Xyleborus bicolor, Blandf., var. α .

Head, legs, and prothorax yellow, the elytra dark reddish brown. *Front* subconvex, sparsely punctured, with short, obscure, median, longitudinal line over the epistoma; the eyes deeply emarginate. *Prothorax* longer than broad, the sides subparallel, slightly broader in front, the basal angles rounded, anteriorly transversely rugose as far as the transverse central nodosity, posteriorly shining and smooth, with a few minute punctures near the base especially. *Scutellum* very minute. *Elytra* the same breadth as the base of the prothorax and one-half longer; lineate-punctate, interstices flat, with uniseriate erect hairs, the sides subparallel, rounded on the apical fifth; viewed laterally, the apical declivity is abrupt, the first and second interstices tuberculate and setose, but not elevated.

Length, 1.7 mm.; breadth, .6 mm.

India: Sunderbans Div. (*C. F. C. Beeson*).

Probably only a local variety of the Japanese species *X. bicolor*, Blandf., from which it differs in the absence of the median longitudinal line on the prothorax, the shorter and more abrupt elytral declivity, and the flatter first and second interstices on the declivity, each of which differences appears to be constant.

XXXIV.—*A Revision of the Genera of the Family Liparidæ.*

By Colonel C. SWINHÖE, M.A., F.L.S., F.Z.S., F.E.S.,
Member of the Entomological Society of France and of
the Bombay Natural History Society.

[Continued from p. 97.]

658. *Aroa incerta*.

Aroa incerta, Rog. Ann. Nat. Hofmus. vi. p. 164 (1891); Kirby, *l. c.* p. 921.

Taveta.

659. *Aroa luisa*.

Aroa luisa, Pag. Jahrb. Nass. Ver. Naturk. xxxviii. p. 26, pl. xi. fig. 3 (1885); Semper, Het. Philipp. ii. p. 456 (1898).

Type, Nias, in Coll. Pagenstecker, Luzon, Philippines.

Ann. & Mag. N. Hist. Ser. 9. Vol. xi.

19

660. *Aroa leucogramma*.

Cænina leucogramma, Felder, Reise Nov. pl. xcix. fig. 7 (1875).

Aroa leucogramma, Semper, *l. c.*

Type, Luzon, in Coll. Rothschild.

661. *Aroa nigrofascia*.

Aroa nigrofascia, Wileman, Entom. 1911, p. 149.

Types, ♂ ♀, Kanshirei, Formosa, in Coll. Wileman.

662. *Aroa globula*.

Aroa globula, Swinhoe, Ann. & Mag. Nat. Hist. (7) xvii. p. 544 (1910).

Type, ♂, Padang, W. Sumatra, in B.M.

663. *Aroa discalis*.

Aroa discalis, Walker, iv. p. 792 (1855), ♂.

Aroa signata, Walker, xxxii. p. 328 (1865), ♀.

Type, ♂, S. Africa; type, ♀, Natal: both in B.M. Mashonaland, Zululand, Port Ogowé, Kilimanjaro, Zomba, Pietermaritzburgh.

664. *Aroa bistigmigera*.

Aroa bistigmigera, Butler, P. Z. S. 1896, p. 847, pl. xlii. fig. 7.

Type, ♂, Nyassaland, in B.M.; Zululand, Port Ogowé, Natal, Basutoland, Nigeria, British East Africa.

665. *Aroa dregei*.

Orgyia dregei, Herr.-Schäff. Aussereur. Schmett. i. fig. 1121 (1854); Kirby, *l. c.* p. 495.

Cape.

666. *Aroa melaxantha*.

Orgyia melaxantha, Walker, xxxii. p. 324 (1865).

Type, ♂, Cape, in B.M.

667. *Aroa socrus*.

Gynæphora socrus, Geyer, Zutr. v. p. 12, figs. 837, 338 (1837).

Aroa socrus, Kirby, *l. c.* p. 464.

Java.

668. *Aroa risoria*.

Aroa risoria, Swinhoe, Trans. Ent. Soc. 1903, p. 457.

Type, ♂, Arjuno, Java, in B.M.

669. *Aroa charax*.

Neurophana charax, Druce, Ann. & Mag. Nat. Hist. (6) xvii. p. 352 (1896).

Type, E. Africa, in Coll. Joicey ; Nyassaland.

670. *Aroa ochraceata*.

Aroa ochraceata, Walker, xxxii. 327 (1865).

Type, ♀, Natal, in B.M. ; Salisbury, Mashonaland, Zomba.

671. *Aroa quadripunctata*.

Orgyia quadripunctata, Wallgrn. Öfv. Vet.-Akad. Forh. xxxii. (1) p. 99 (1876).

Transvaal, Bine, W. Africa.

672. *Aroa niasana*.

Aroa niasana, Swinhoe, Ann. & Mag. Nat. Hist. (7) xix. p. 203 (1907).

Type, ♂, Nias, in B.M.

673. *Aroa flaveofusca*.

Aroa flaveofusca, Swinhoe, l. c. (7) ix. p. 80 (1902).

Type, ♂, Lawas, Borneo, in B.M.

674. *Aroa mirma*.

Aroa mirma, Swinhoe, Trans. Ent. Soc. 1903, p. 455.

Type, ♂, Sumatra, in B.M.

675. *Aroa plana*.

Orgyia (?) *plana*, Walker, iv. p. 786 (1855).

Aroa plana, Strand, l. c. pl. xxii. d.

Charnidas junctifera, Walker, xxxii. p. 334 (1865).

Type, ♂, N. India, in B.M. ; type, *junctifera*, in Layard's lost collection.

676. *Aroa substrigosa*.

Aroa substrigosa, Walker, iv. p. 794 ; Butler, Ill. Het. v. p. 54, pl. xc. fig. 5 (1886) ; Strand, l. c. pl. xix. b.

Type, ♂, Sylhet, in B.M. ; Chin Hills, Khasia Hills.

677. *Aroa pyrrhochronia*.

Aroa pyrrhochronia, Walker, xxxii. p. 329 (1865) ; Strand, l. c. pl. xix. i.

Buziza detecta, Walker, xxxii. p. 399.

Type, ♂, Cherapunji, in B.M. ; Mandi Bukar, Bhootan, Sikkim, Jaintia Hills.

Hampson, in 'Moths of India,' puts these two and *clara* as one species, *substrigosa* and *pyrrhochroma* may be of one very variable species, but I have a fine series of both; *clara* is undoubtedly quite distinct, being of a different colour, pale yellowish fawn-colour. I have one male and three females in my collection.

678. *Aroa clara*.

Aroa clara, Swinhoe, *l. c.* fig. 10.

Type, Bombay, in B.M.; Poona.

679. *Aroa cinnamomea*.

Charnidas cinnamomea, Moore, Lep. Atk. p. 44 (1879).

Charnidas aurantiaca, Warren, P. Z. S. 1888, p. 296.

Type, ♂, Kunawar; type, ♂, *aurantiaca*, Attock: both in B.M.

680. *Aroa flavicollis*.

Crinola flavicollis, Leech, Entom. xxiii. p. 111 (1896), and P. Z. S. 1899, p. 120.

Aroa flavicollis, Strand, *l. c.* p. 121, pl. xxii. c.

Form *leucoides*, Strand, *l. c.*

Type, ♂, Chang-Yang; type, ♀, Chia-kow: both in B.M.

681. *Aroa callima*.

Aroa callima, Beth.-Baker, *l. c.* (8) vii. p. 546.

Type, ♀, Idakun, Lagos, in Coll. Bethune-Baker.

682. *Aroa perfida*.

Aroa perfida, Beth.-Baker, *l. c.*

Type, ♂, Gunnal, W. Africa, in Coll. Bethune-Baker.

683. *Aroa catori*.

Aroa catori, Beth.-Baker, *l. c.* p. 546.

Type, ♂, Lokoja District, in Coll. Bethune-Baker.

684. *Aroa pinodes*.

Aroa pinodes, Beth.-Baker, *l. c.* p. 547.

Type, ♂, N'Dalla Tondo, in Coll. Bethune-Baker.

685. *Aroa obliqua*.

Aroa obliqua, Beth.-Baker, *l. c.* p. 546.

Type, ♂, Lokoja District, in Coll. Bethune-Baker.

Genus LACHANA, Moore, P. Z. S. 1888, p. 397.

686. *Lachana ladakensis*.

Lachana ladakensis, Moore, *l. c.* p. 398; Butler, Ill. Het. vii. p. 30, pl. cxxi. fig. 6 (1889).

Type, ♂, Ladak, in B.M.; Skardo, Kashmir.

Genus ICTA, Walker, iv. p. 922 (1855).

687. *Icta fulviceps*.

Icta fulviceps, Walker, *l. c.*

Type, ♂, Australia, in B.M.

688. *Icta tanaopis*.

Icta tanaopis, Turner, Proc. Linn. Soc. N.S.W. xlv. (4) p. 477 (1920).

Type, Claudie River, N. Queensland, in Coll. Turner.

Genus VARMINA, Moore, P. Z. S. 1888, p. 405.

689. *Varmina indica*.

Gluphisia (?) *indica*, Walker, v. p. 1039 (1855).

Type, India, in Mus. Oxon.; Simla, Kangra, Dharmsala, Umballa.

Genus DASORGYIA, Staudinger, Stett. ent. Zeit. xlii.
p. 406 (1881).

690. *Dasorgyia pumila*.

Dasychira pumila, Staud. *l. c.* p. 605.

Saisan, Tarbagatai District.

691. *Dasorgyia selenophora*.

Dasorgyia selenophora, Staud. *l. c.* xlviii. p. 98 (1887).

Var. *pamirica*, Strand, *l. c.* pl. xxii. *b.*

Ferghana, Pamir.

692. *Dasorgyia alpherakii*.

Dasorgyia alpherakii, Gr.-Grsh. Hor. Soc. Ent. Ross. xxv. p. 464 (1891); Strand, *l. c.* pl. xxii. *b.*

Kuku-Nor and Amdo District.

693. *Dasorgyia grumi*.

Dasychira grumi, Staud. Cat. Pal. Lep. p. 115.

Dasychira semenovi, Gr.-Grsh. *l. c.*

Kuku-Nor (Tibet).

Genus *CASAMA*, Walker, xxxiii. p. 611 (1865).

694. *Casama vilis*.

Eproctis vilis, Walker, xxxii. p. 348, ♂.

Casama indeterminata, Walker, xxxii. p. 611, ♀.

Types, ♂ ♀, S. India, in B.M.; Nilgiris, Ceylon, Hyderabad, Sind, Muscat, Arabia, Somaliland, Abyssinia.

695. *Casama innotata*.

Spilosoma innotata, Walker, iii. p. 674 (1885).

Sierarctia innotata, Kirby, l. c. p. 243.

Egypt.

696. *Casama uniformis*.

Oeneria uniformis, Roths. Nov. Zool. xx. p. 118 (1913).

Casama uniformis, Roths. l. c. xxiv. p. 359 (1917).

Type, Algeria, in Coll. Rothschild.

Genus *THIACIDAS*, Walker, v. p. 1027 (1856).

697. *Thiacidas postica*.

Thiacidas postica, Walker, v. p. 1028, ♂.

Drymonia (?) *denotata*, Walker, xxxii. p. 414, ♀.

Heterocampa (?) *nigroscripta*, Walker, xxxii. p. 423, ♀.

Cnethocampa (?) *curvata*, Walker, xxxii. p. 429, ♂.

Cnethocampa (?) *basifurca*, Walker, xxxii. p. 430, ♂.

Type, Nepal; types of all the others, India: all in B.M. Poona, Bombay, Burma.

Genus *IROPOCA*, Turner, Trans. Ent. Soc. 1904, p. 477.

698. *Iropoca rotundata*.

Teara rotundata, Walker, iv. p. 851.

Iropoca rotundata, Turner, Proc. Roy. Soc. N.S.W. xlv. (4) p. 494 (1920).

Type, ♂, Tasmania, in B.M.; Moreton Bay, Brisbane; Jervis Bay, near Nowra; Beechworth; Gisborne.

Genus *AXIOLOGA*, Turner, Trans. Ent. Soc. 1904, p. 472.

699. *Axiologa pura*.

Teara pura, Lucas, Proc. Roy. Soc. Queensland, 1891, p. 75.

Type, Queensland, in Coll. Lucas.

Genus CADUCA, Swinhoe, Ann. & Mag. Nat. Hist. (7) xvii.
p. 546 (1906).

700. *Caduca venata*.

Caduca venata, Swinhoe, *l. c.*

Type, ♂, Bihé, W. Africa, in B.M.

Genus ORGYIA, Ochs. Schmett. Eur. ii. p. 208 (1810);
type, *pudibunda*, Linn.

Dasychira, Steph. Ill. Brit. Ent. Haust. ii. p. 58 (1829); type, *pudibunda*, Linn.

Psalis, Hübner, Zutr. ii. p. 19 (1823); type, *securis*, Hübner.

Olene, Hübner, *l. c.*; type, *mendosa*, Hübner.

Arestha, Walker, iv. p. 805 (1855); type, *antica*, Walker.

Melia, Walker, iv. p. 808; type, *costalis*, Walker.

Anaxila, Walker, iv. p. 810; type, *plagiata*, Walker.

Argila, Walker, iv. p. 811; type, *basalis*, Walker.

Niocha, Walker, v. p. 1069 (1855); type, *fusiformis*, Walker.

Dediana, Walker, v. p. 1074; type, *antica*, Walker.

Rilia, Walker, v. p. 1025; type, *lunceolata*, Walker.

Phineca, Walker, vii. p. 1745 (1856); type, *basistriga*, Walker.

Iiema, Moore, Cat. Lep. E. I. Co. ii. p. 341 (1859); type, *costalis*, Walker.

Anchyneura, Felder, Sitz. Akad. Wiss. Wien, xliii. (i.) p. 33 (1861); type, *præusta*, Felder.

Thelda, Walker, Journ. Linn. Soc. vi. p. 139 (1862); type, *patula*, Walker.

Boreconia, Walker, xxxii. p. 460 (1865); type, *subviridis*, Walker.

Tearosoma, Felder, Reise Nov. pl. c. fig. 6 (1868); type, *aspersum*, Felder.

Twiriga, Walker, Char. Undescr. Lep. Het. p. 15 (1869); type, *invasa*, Walker.

Cadrusia, Moore, Lep. Atk. p. 54 (1879); type, *virescens*, Moore.

Calliteara, Butler, Trans. Ent. Soc. 1881, p. 12; type, *argentata*, Butler.

Eudasychira, Mösch. Abh. Senck. Ges. xv. p. 75 (1889); type, *quinquepunctata*, Mösch.

Pseudonotodonta, Mösch. *l. c.* p. 76; type, *virescens*, Mösch.

Æcura, Holland, Psyche, vi. p. 339 (1893); type, *goodii*, Holland.

Euproctidion, Holland, *l. c.* p. 414; type, *gabunica*, Holland.

Notohyba, Holland, *l. c.* p. 434; type, *striata*, Holland.

Thannocera, Holland, *l. c.* p. 454; type, *albinotata*, Holland.

Bathmochtha, Karsch, Ent. Nachr. xxi. p. 368 (1895); type, *albilunulata*, Karsch.

701. *Orgyia pudibunda*.

Bombyx pudibunda, Linn. Syst. Nat. x. p. 363 (1758).

Dasychira pudibunda, Leech, Trans. Ent. Soc. 1899, p. 124; Strand, *l. c.* pl. xix. d, ♀.

Geometra scopularia, Linn. *l. c.* i. p. 521 (1758).

Noctua justica, Müll. Faun. Fridr. p. 47 (1764).

Phalena porrecta, Retz. Gen. Spec. Ins. p. 38 (1783).

Bombyx juglandis, Hübner, Eur. Schmett. Bomb. figs. 84, 85 (1800).

Dasychira pudibunda, var. *concolor*, Staud. Cat. Lep. Eur. p. 29 (1861);
Strand, l. c. pl. xix. e.

S. Europe; Armenia; Syria; Amurland; Ussuri; N.E. China; Japan.

702. *Orgyia olga*.

Dasychira olga, Oberth. Etude d'Ent. v. p. 34, pl. ii. figs. 1, 2 (1881);
Strand, l. c. p. 113, pl. xix. g.

S.E. Siberia.

703. *Orgyia fascellina*.

Bombyx fascellina, Linn. Syst. Nat. i. p. 503 (1758).

Dasychira fascellina, Strand, l. c. pl. xix. f.

Bombyx medicaginis, Hübner, Eur. Schmett. ii. fig. 81 (1800).

Dasychira laricis, Schille, Soc. Ent. xiv. p. 403.

Orgyia obscura, Zeller, Ins. Lep. p. 927 (1809).

Dasychira unicolor, Schultz, Entom. Zeit. xxiv. p. 35.

Dasychira nivalis, Staud. Stett. Zg. 1887, p. 97.

Dasychira angelus, Tschelv. Rev. Russe. Ent. iv. p. 77.

Dasychira proletaria, Strand, l. c.

Europe; Altai, Mongolia, Vallais; Alai, Transalai, Samarkand, Tian Shan.

704. *Orgyia fortunata*.

Dasychira fortunata, Rog. Vehr. zool.-bot. Ges. Wien, 1891, p. 566;
Strand, l. c. pl. xix. g.

Canary Islands.

705. *Orgyia abietis*.

Bombyx abietis, Schiff. Syst. Verz. Wien, p. 55 (1776).

Dasychira mediobscura, Schultz, Entom. Zachr. Stuttgr. xxiv. p. 35.

Dasychira albietis, Strand, l. c. pl. xix. g.

Sweden; Russia; Germany; Carpathians; Bohemia; Austria.

706. *Orgyia pseudabietis*.

Calliteara pseudabietis, Butler, Cist. Ent. iii. p. 118 (1885).

Calliteara abietis, Leech (nec Schiff.), P. Z. S. 1888, p. 631.

Dasychira pryer, Butler, l. c. p. 119.

Dasychira pulica, Staud. Rom. sur Léop. iii. p. 204 (1887).

Dasychira modesta, Kirby, Cat. Moths, i. p. 483 (1892).

Dasychira pseudabietis, Strand, l. c. pl. xix. h.

Types, ♂ ♀, Japan, in B.M.; Amur.

707. *Orgyia argentata*.

Dasychira argentata, Butler, Trans. Ent. Soc. 1881, p. 12.

Type, Japan, in B.M.

708. *Orgyia selenitica*.

Bombyx selenitica, Esper, Schmett. iii. (2) p. 18, pl. lxxxii. fig. 5, p. 49, pl. lxxxviii. fig. 2 (1783).

Bombyx paradoxa, Fabr. Mant. Ins. ii. p. 130 (1787).

Bombyx lathyræ, Hübner, Eur. Schmett. Bomb. figs. 79, 80 (1800).

Dasychira selenitica, Strand, l. c. pl. xix. b.

Germany; Austria; Russia; Finland.

709. *Orgyia lunulata*.

Dasychira lunulata, Butler, Ann. & Mag. Nat. Hist. (4) xx. p. 403 (1877); Butler, Ill. Het. ii. pl. xxiv. fig. 8 (1878).

Dasychira solitaria, Staud. Rom. sur Léop. iii. pl. xii. fig. 1 (1878).

Dasychira acronycta, Oberth. Etud. d'Ent. v. p. 35, pl. v. fig. 7, ♂ (1881).

Type, Japan, in B.M.; Askold, Vladivostock.

710. *Orgyia virginia*.

Dasychira virginia, Oberth. Diagn. Lep. Askold, p. 7 (1879); Oberth. Etud. d'Ent. p. 33, pl. v. fig. 5 (1881); Wileman, Trans. Ent. Soc. 1911, p. 271.

Askold, near Vladivostock; Japan.

711. *Orgyia dalbergiæ*.

Dasychira dalbergiæ, Moore, P. Z. S. 1888, p. 399; Strand, l. c. pl. xxii. a.

Dasychira kangrana, Strand, l. c. p. 115.

Type, Dharmasala, Khasia Hills.

712. *Orgyia invaria*.

Repena (?) *invaria*, Walker, vii. p. 1724 (1856).

Dasychira inclusa, Walker, vii. p. 1727.

Thelde patula, Walker, Journ. Linn. Soc. vi. p. 140 (1862).

Type, ♂, Philippines; type, ♂, *inclusa*, Java: both in B.M.

Type, *patula*, Sarawak, in Mus. Oxon.; Sumatra, Sumbawa, Hongkong.

713. *Orgyia juncta*.

Dasychira juncta, Wileman, Trans. Ent. Soc. 1911, p. 270; Wileman, The Philip. Journ. of Science, xiii. (4) p. 151, pl. i. figs. 1, 2 (1918).

Type, ♂, Yoshima Yama, Japan, in Coll. Wileman.

714. *Orgyia horsfieldii*.

Arctia horsfieldii, Moore, Trans. Ent. Soc. 1851, p. 162, pl. xii. figs. 1-3.

Dasychira horsfieldii, Swinhoe, Trans. Ent. Soc. 1903, p. 473 (note);

Turner, Proc. Linn. Soc. N.S.W. xlv. (4) p. 497 (1920).

Dasychira arga, Moore, Cat. Lep. E. I. Co. ii. p. 339 (1859).

Dasychira grotei, Moore, *l. c.* p. 338; Strand, *l. c.* p. 115, pl. xxii. *a*.

Dasychira varia (♀ only), Walker, iv. 868 (1855).

Daschira longipennis, Walker, Journ. Linn. Soc. vi. p. 13 (1862).

Dasychira kansalia, Moore, P. Z. S. 1879, p. 401.

Dasychira nilgirica, Hmps. Ill. Het. viii. p. 58, pl. cxiv. figs. 13, 14 (1891).

Teara farenooides, Lucas, Proc. Roy. Soc. Queensland, 1892, p. 75.

Types *horsfieldii* and *longipennis* should be in Mus. Oxon., but I have been unable to find them. Type, ♀, *grotei*, N. India; type, ♂, *kansalia*, Kasaoli; types, ♂ ♀, *nilgirica*, Nilgiris; types, ♂ ♀, *arga*, Java: all in B.M. Singapore and many parts of India and Ceylon, also in Australia, Cairns, Brisbane.

715. *Orgyia flavimacula*.

Dasychira flavimacula, Moore, P. Z. S. 1865, p. 804; Swinhoe, Trans.

Ent. Soc. 1903, p. 465 (note).

Form *yatonga*, Strand, *l. c.* p. 116.

Type, ♂, Bengal, in B.M.; Sikkim; Yatung, Tibet.

716. *Orgyia securis*.

Psalis securis, Hübner, Zutr. iii. p. 9, figs. 291, 292 (1823).

Dasychira securis, Strand, *l. c.* pl. xxii. *c*.

Aresta antica, Walker, iv. p. 805 (1855).

Rigema falcata, Walker, xxxii. p. 437 (1865).

Rigema tacta, Walker, *l. c.* p. 438.

Anticyra approximata, Walker, xxxii. p. 440.

Type, ♂, *antica*, Java; type, ♂, *falcata*, Ceylon; type, *tacta*, Australia; type, *approximata*: all in B.M. Australia, Cairns, Rockhampton, Duaringa, Brisbane, Philippines, and many parts of India and Africa.

717. *Orgyia olearia*.

Olene olearia, Swinhoe, P. Z. S. 1885, p. 297, pl. xx. figs. 14, 15.

Types, ♂ ♀, Poona, in B.M.; Nilgiris; Madras; Ahmednagar.

718. *Orgyia fusca*.

Anthora fusca, Walker, iv. 918 (1855), ♂.

Bareconia subviridis, Walker, xxxii. p. 460 (1865), ♀ (nec ♂).

Types, ♂ ♀, S. Africa, in B.M.; Cape; Knysna.

719. *Orgyia albiplaga*.

Dasychira albiplaga, Swinhoe, Ann. & Mag. Nat. Hist. (8) i. p. 62 (1908).

Type, ♀, E. Java, in B.M.; Talaut Island.

720. *Orgyia cæruleifascia*.

Thamnocera cæruleifascia, Holland, Pysche, vi. p. 469 (1893).

Type, ♀, Ogove River, in Coll. Holland; River Niger.

721. *Orgyia robusta*.

Acyphas (?) *robusta*, Walker, iv. p. 799.

Type, ♂, Sierra Leone, in B.M.

722. *Orgyia delicata*.

Notohyba delicata, Holland, l. c. p. 451, pl. xviii. fig. 4.

Type, ♀, Ogove River, in Coll. Holland.

723. *Orgyia citana*.

Utidea (?) *citana*, Schaus & Clements, Sierra Leone Lep. p. 26, pl. i. fig. 10 (1893).

Sierra Leone.

724. *Orgyia punctifera*.

Erastris (?) *punctifera*, Walker, xii. 809 (1857).

Type, ♂, Congo, in B.M.

725. *Orgyia peculiaris*.

Mardara peculiaris, Butler, Ann. & Mag. Nat. Hist. (8) iv. p. 240 (1879).

Type, ♂, Madagascar, in B.M.

726. *Orgyia basistriga*.

Phineca basistriga, Walker, vii. p. 1747 (1856).

Type, ♀, without locality, in B.M.

727. *Orgyia antica*.

Dediana antica, Walker, v. p. 1074 (1855).

Bathmochtha albilunulata, Karsch, Ent. Nachr. xxi. p. 368, pl. iv. fig. 12 (1895).

Type, ♂, without locality, in B.M.; type, *albilunulata*, Cameroons.

728. *Orgyia lignea*.

Nioda lignea, Butler, Ann. & Mag. Nat. Hist. (5) iv. p. 241 (1879).

Type, ♂, Madagascar, in B.M.

729. *Orgyia basalis*.

Argila basalis, Walker, iv. p. 811.

Type, ♂, Sierra Leone, in B.M.

730. *Orgyia quinquepunctata*.

Eudasychira quinquepunctata, Mösch. Abhandl. Senck. Nat. Ges. xv. p. 75 (1887); Kirby, l. c. p. 485.

Accra.

731. *Orgyia lignifica*.

Rilia (?) *lignifica*, Walker, Entom. v. p. 125 (1869).

Olene (?) *lignifica*, Kirby, l. c. p. 488 (1892).

Red Sea.

732. *Orgyia basifurca*.

Anthora basifurca, Walker, xxxii. p. 335 (1865).

Argila basifurca, Kirby, l. c. p. 491.

Type, Caffraria, in B.M.

733. *Orgyia basigutta*.

Phineca basigutta, Walker, vii. p. 1746 (1856).

Habitat. ignotus, type, in B.M.

734. *Orgyia orimba*.

Olene orimba, Swinhoe, Ann. & Mag. Nat. Hist. (6) xiv. p. 435 (1894).

Types, ♂ ♀, Khasia Hills, in B.M.

735. *Orgyia anana*.

Dasychira anana, Swinhoe, l. c. (7) xviii. p. 406 (1910).

Type, Java, in B.M.; Sumatra.

736. *Orgyia pulchra*.

Dasychira pulchra, Swinhoe, l. c. (7) xvii. p. 544 (1906).

Type, ♂, Coomassie, in B.M.

737. *Orgyia postalba*.

Dasychira postalba, Swinhoe, l. c. p. 545.

Type, ♂, Coomassie, in B.M.

738. *Orgyia kanshireiensis*.

Psalis (?) *kanshireiensis*, Wileman, Entom. 1917, p. 146.

Type, ♂, Kanshirei, Formosa, in Coll. Wileman.

739. *Orgyia transversa*.

Nioda transversa, Walker, xxxiv. p. 434 (1866).

Type, Ceylon, in B.M.

740. *Orgyia hieroglyphica*.

Dasychira hieroglyphica, Swinhoe, Trans. Ent. Soc. 1904, p. 144.

Type, ♀, Obuassi, Ashanti, in B.M.

741. *Orgyia præusta*.

Anchyneura præusta, Felder, Sitz. Akad. Wiss. Wien, xliii. (i.) p. 33 (1861).

Arestha præusta, Felder, Reise. Nov. pl. xcix. fig. 5 (1868).

Type, Amboina, in Coll. Rothschild.

742. *Orgyia davidi*.

Nerice davidi, Oberth. Etud. d'Ent. vi. p. 17, pl. ix. fig. 2 (1881); Kirby, Cat. Moths, i. p. 487 (1892).

China.

743. *Orgyia bipartita*.

Nerice bipartita, Butler, Cist. Ent. iii. p. 119 (1885); Kirby, *l. c.*

Type, Japan, in B.M.

744. *Orgyia asvata*.

Dasychira asvata, Moore, Cat. Lep. E. I. Co. ii. p. 340 (1859).

Types, ♂ ♀, Java, in B.M.; Dinding Isl., Malaya.

745. *Orgyia rendalli*.

Dasychira rendalli, Distant, Ann. & Mag. Nat. Hist. (5) xx. p. 203 (1897).

Type, Transvaal, in Coll. Distant; Natal; Mashonaland.

746. *Orgyia municipalis*.

Lælia municipalis, Distant, *l. c.* p. 200, ♂.

Lacipa diffusa, Distant, *l. c.*, ♀.

Types, ♂ ♀, Transvaal, in Coll. Distant; Cape; Natal; Nyassaland.

747. *Orgyia extorta*.

Dasychira extorta, Distant, *l. c.* p. 203.

Lymantria hera, Druce, Ann. & Mag. Nat. Hist. (7) i. p. 209 (1898).

Type, Transvaal, in Coll. Distant; type, *hera*, Dar-es-Salaam, in Coll. Joicey; Natal; Mashonaland.

748. *Orgyia nubifuga*.

Dasychira nubifuga, Holland, Psyche, vi. p. 451, pl. xviii. fig. 33 (1893).

Type, ♂, Ogove River, in Coll. Holland.

749. *Orgyia miserata*.

Ilema miserata, Holland, *l. c.* pl. xvii. fig. 3.

Type, ♂, Ogove River, in Coll. Holland; Lokoja.

750. *Orgyia ocellifera*.

Æcura ocellifera, Holland, *l. c.* p. 454, pl. xviii. fig. 5.

Dasychira ocellifera, Swinhoe, Trans. Ent. Soc. 1903, p. 466 (note).

Type, ♂, Ogove River, in Coll. Holland.

751. *Orgyia glovera*.

Dasychira glovera, Swinhoe, Ann. & Mag. Nat. Hist. (7) xvii. p. 545 (1906).

Type, ♂, Entebbe, Uganda, in B.M.

752. *Orgyia rocana*.

Dasychira rocana, Swinhoe, *l. c.*

Type, ♂, Coomassie, in B.M.

753. *Orgyia pytna*.

Dasychira pytna, Druce, Ann. & Mag. Nat. Hist. (7) i. p. 210 (1898).

Type, ♂, Gambia, W. Africa, in Coll. Joicey.

754. *Orgyia maligna*.

Parorgyia maligna, Butler, Cist. Ent. iii. p. 17 (1882).

Type, ♂, Madagascar, in B.M.

755. *Orgyia phasiana*.

Parorgyia phasiana, Butler, *l. c.*

Type, ♂, Madagascar, in B.M.

756. *Orgyia chalana*.

Dasychira chalana, Moore, Cat. Lep. E. I. Co. ii. p. 339 (1859).

Lacida costiplaga, Walker, Journ. Linn. Soc. vi. p. 126 (1862).

Type, ♀, Java, in B.M.; type, ♂, *costiplaga*, Sarawak, in Mus. Oxon.

757. *Orgyia misana*.

Dasychira misana, Moore, *l. c.* p. 340, pl. ii. a, fig. 2; Snellen, Tijds. v. Ent. xxii. p. 110 (1879).

Type, ♂, Java, in B.M.; Celebes.

758. *Orgyia ostracina*.

Lælia ostracina, Turner, Trans. Roy. Soc. S. Australia, 1902, p. 181.

Dasychira ostracina, Turner, Proc. Linn. Soc. N.S.W. xlv. (4) p. 497 (1920) (note).

Cape York.

759. *Orgyia whitei*.

Æcura whitei, Druce, Ann. & Mag. Nat. Hist. (7) i. p. 209 (1898).

Type, ♂, Old Calabar, in Coll. Joicey; Natal; Durban.

760. *Orgyia goodii*.

Æcura goodii, Holland, Psyche, vi. p. 390, pl. vii. figs. 3, 4 (1893).

Type, ♂, Ogove River, in Coll. Holland; Sierra Leone.

761. *Orgyia libella*.

Dasychira libella, Swinhoe, Trans. Ent. Soc. 1904, p. 145.

Type, ♂, Obuassi, Ashanti, in B.M.

762. *Orgyia costalis*.

Melia costalis, Walker, iv. p. 808 (1855).

Type, ♀, Java, in B.M.; Heipaw, Burma.

763. *Orgyia cuneifera*.

Melia cuneifera, Walker, Journ. Linn. Soc. vi. p. 127 (1862).

Type, Sarawak, in B.M.

764. *Orgyia cromptoni*.

Dasychira cromptoni, Swinhoe, Trans. Ent. Soc. 1903, p. 467 (note).

Type, ♂, Old Calabar, in B.M.; River Niger.

765. *Orgyia anaha*.

Dasychira anaha, Swinhoe, Ann. & Mag. Nat. Hist. (7) xviii. p. 406 (1906).

Type, ♂, Padang, Sumatra; type, ♀, Batavia, Java: both in B.M.

766. *Orgyia bhana*.

Dasychira bhana, Moore, P. Z. S. 1865, p. 804; Swinhoe, Trans. Ent. Soc. 1903, p. 468 (note).

Type, ♂, Darjiling, in B.M.

767. *Orgyia nigracula*.

Dasychira nigracula, Walker, xxxii. p. 360 (1865); Swinhoe, *l. c.* (note).

Type, ♂, N. India, in B.M.; Sikkim, Khasia Hills.

768. *Orgyia feminula*.

Mardara feminula, Hmps. Ill. Het. viii. p. 58, pl. cxli. fig. 1-7 (1891).
Dasychira feminula, Swinhoe, *l. c.* (note).

Types, ♂ ♀, Nilgiri Hills, in B.M.

769. *Orgyia magnalia*.

Dasychira magnalia, Swinhoe, Ann. & Mag. Nat. Hist. (7) xii. p. 198 (1903).

Types, ♂ ♀, Khasia Hills, in B.M.

770. *Orgyia cerebosa*.

Lymantria cerebosa, Swinhoe, Trans. Ent. Soc. 1903, p. 483.

Type, ♂, Solon, Simla, in B.M.

771. *Orgyia chloroptera*.

Dasychira chloroptera, Hmps. Moths of India, i. p. 450 (1880).

Type, ♂, Khasia Hills, in B.M.

772. *Orgyia angiana*.

Dasychira angiana, Joicey, Noakes, & Talbot, Trans. Ent. Soc. 1915, p. 382, pl. lxii. fig. 1.

Type, ♂, Arfak Mts., Dutch N. Guinea, in Coll. Joicey.

773. *Orgyia nigra*.

Dasychira nigra, Hmps. Journ. Bombay Nat. Hist. Soc. xiii. p. 416, pl. ii. fig. 19 (1900); Wileman, Trans. Ent. Soc. 1911, p. 270.

Type, ♀, Sikkim, in B.M.; Japan.

[To be continued.]